

09-25-00

A

Please type a plus sign (+) inside this box → ☐

PTO/SB/05 (4/98)

Approved for use through 09/30/2000. OMB 0651-0032
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 C.F.R. § 1.53 (b))

Attorney Docket No.	LEX-0042-USA
First Inventor or Application Identifier	Gregory Donoho et al.
Title	Novel Human Protease Inhibitor-Like Proteins and Polynucleotides Encoding the Same
Express Mail label No.	EL584856694US

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents

ADDRESS TO: Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

1	<input type="checkbox"/>	*Fee Transmittal Form (e.g., PTO/SB/17) (Submit an original and a duplicate for fee processing)	
2	<input checked="" type="checkbox"/>	Specification [Total 25] Pages (preferred arrangement set forth below)	
<ul style="list-style-type: none">- Descriptive title of the invention- Cross References to Related Applications- Statement Regarding Fed sponsored R & D- Reference to Microfiche Appendix- Background of the invention- Brief Summary of the invention- Brief Description of the drawings (if filed)- Detailed Description- Claim(s)- Abstract of the disclosure			
3	<input type="checkbox"/>	Drawing(s) (35 U.S.C. 113) [Total 1] Sheets	
4	<input type="checkbox"/>	Oath or Declaration [Total 1]	
a. <input checked="" type="checkbox"/> Newly unexecuted (original or copy)			
b. <input type="checkbox"/> Copy from a prior application (37 C.F.R. § 1.63(d)) (for continuation/divisional with Box 16 completed)			
i. <input type="checkbox"/> DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).			

5	<input type="checkbox"/>	Microfiche Computer Program (Appendix)
6	<input type="checkbox"/>	Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
a. <input type="checkbox"/> Computer Readable Copy		
b. <input checked="" type="checkbox"/> Paper Copy (identical to computer copy)		
c. <input type="checkbox"/> Statement verifying identity of above copies		

ACCOMPANYING APPLICATION PARTS

7	<input type="checkbox"/>	Assignment Papers (cover sheet & document(s))
8	<input type="checkbox"/>	37 C.F.R. § 3.73(b) Statement (when there is an assignee) <input type="checkbox"/> Power of Attorney
9	<input type="checkbox"/>	English Translation Document (if applicable)
10	<input type="checkbox"/>	Information Disclosure Statement (IDS)/PTO-1449 <input type="checkbox"/> Copies of IDS Citations
11	<input type="checkbox"/>	Preliminary Amendment
12	<input checked="" type="checkbox"/>	Return Receipt Postcard (MPEP 503) (Should be specifically itemized)
13	<input checked="" type="checkbox"/>	*Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application, Status still proper and desired (PTO/B/09-12)
14	<input type="checkbox"/>	Certified Copy of Priority Document(s) (if foreign priority is claimed)
15	<input type="checkbox"/>	Other:

*NOTE FOR ITEMS 1 & 13: IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELED UPON (37 C.F.R. § 1.28).

16. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:

<input type="checkbox"/> Continuation	<input type="checkbox"/> Divisional	<input type="checkbox"/> Continuation-in-part (CIP) of prior application No: _____
---------------------------------------	-------------------------------------	--

Prior application information: Examiner: _____ Group/Art Unit: _____

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference.

17. CORRESPONDENCE ADDRESS

<input checked="" type="checkbox"/> Customer Number or Bar Code Label	(Insert Customer No. or Attach bar code label here)	or <input type="checkbox"/> Correspondence address below
Name	24231 PATENT TRADEMARK OFFICE	
Address		
City	State	Zip Code
Country	Telephone (281) 362-6554	Fax (281) 364-0155

Name (Print/Type)	Lance K. Ishimoto	Registration No.	41866
Signature		Date	Sept. 22, 2000

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

**STATEMENT CLAIMING SMALL ENTITY STATUS
(37 CFR 1.9(f) & 1.27(c)) -- SMALL BUSINESS CONCERN**

Docket Number (Optional)
LEX-0042-USA

Applicant, Patentee, or Identifier: Gregory Donoho et al.
Application or Patent No.: _____
Filed or Issued: September 22, 2000
Title: Novel Human Protease Inhibitor-Like Proteins and Polynucleotides Encoding the Same

I hereby state that I am

- ☐ the owner of the small business concern identified below:
☒ an official of the small business concern empowered to act on behalf of the concern identified below:

NAME OF SMALL BUSINESS CONCERN Lexicon Genetics Incorporated

ADDRESS OF SMALL BUSINESS CONCERN 4000 Research Forest Drive, The Woodlands, TX 77381

I hereby state that the above identified small business concern qualifies as a small business concern as defined in 13 CFR Part 121 for purposes of paying reduced fees to the United States Patent and Trademark Office. Questions related to size standards for a small business concern may be directed to: Small Business Administration, Size Standards Staff, 409 Third Street, SW, Washington, DC 20416.

I hereby state that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention described in:

- ☒ the specification filed herewith with title as listed above.
☐ the application identified above.
☐ the patent identified above.

If the rights held by the above identified small business concern are not exclusive, each individual, concern, or organization having rights in the invention must file separate statements as to their status as small entities, and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 CFR 1.9(e).

Each person, concern, or organization having any rights in the invention is listed below:

- ☒ No such persons, concerns, or organizations exist.
☐ each such person, concern, or organization is listed below:


Separate statements are required from each named person, concern, or organization having rights to the invention stating their status as small entities. (37 CFR 1.27)

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

NAME OF PERSON SIGNING Lance K. Ishimoto

TITLE OF PERSON IF OTHER THAN OWNER Vice President - Intellectual Property

ADDRESS OF PERSON SIGNING 4000 Research Forest Drive, The Woodlands, TX 77381

SIGNATURE  DATE September 22, 2000
Reg. No. 41,866

**NOVEL HUMAN PROTEASE INHIBITOR-LIKE PROTEINS AND POLYNUCLEOTIDES
ENCODING THE SAME**

1. INTRODUCTION

5 The present application claims priority to U. S. Provisional Application Number 60/156,101 which was filed September 24, 1999 which is herein incorporated by reference in its entirety.

10 The present invention relates to the discovery, identification, and characterization of novel human polynucleotides encoding proteins that share sequence similarity with mammalian trypsin inhibitors. The invention encompasses the described polynucleotides, host cell expression systems, the encoded proteins, fusion proteins, polypeptides and peptides, antibodies to the encoded proteins and peptides, and genetically
15 engineered animals that either lack or over express the disclosed genes, antagonists and agonists of the proteins, and other compounds that modulate the expression or activity of the proteins encoded by the disclosed genes that can be used for diagnosis, drug screening, clinical trial monitoring, the treatment of
20 physiological disorders, or otherwise contributing to the quality of life.

2. BACKGROUND OF THE INVENTION

25 Proteases are enzymes that mediate the proteolytic cleavage of polypeptide sequences. Conversely, protease inhibitors prevent or hinder proteolytic activity. Given the importance of proteolysis in a wide variety of cellular functions and disease, protease inhibitors have been demonstrated to be involved in, *inter alia*, regulating development, modulating cellular processes, and preventing infectious, and particularly viral, disease.

3. SUMMARY OF THE INVENTION

30 The present invention relates to the discovery, identification, and characterization of nucleotides that encode

novel human proteins, and the corresponding amino acid sequences of these proteins. The novel human proteins (NHPs) described for the first time herein share structural similarity with animal trypsin inhibitor proteins. As such, the novel genes represent a new class of proteins with a range of homologues and orthologs that transcend phyla and a range of species.

The novel human nucleic acid sequences described herein, encode proteins/open reading frames (ORFs) of 497 amino acids in length (see SEQ ID NO: 2).

The invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof that compete with native NHP, peptides, and antibodies, as well as nucleotide sequences that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and gene or regulatory sequence replacement constructs) or to enhance the expression of the described NHP genes (e.g., expression constructs that place the described gene under the control of a strong promoter system), and transgenic animals that express a NHP transgene, or "knock-outs" (which can be conditional) that do not express a functional NHP.

Further, the present invention also relates to processes of identifying compounds that modulate, i.e., act as agonists or antagonists, of NHP expression and/or NHP product activity that utilize purified preparations of the described NHPs and/or NHP product, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

The Sequence Listing provides the sequences of a trypsin inhibitor-like ORF that encodes the described NHP amino acid sequences.

5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs, described for the first time herein, are novel proteins that are expressed in, *inter alia*, human cell lines, and human prostate, fetal brain, cerebellum, spinal cord, thymus, spleen, lymph node, bone marrow, trachea, lung, kidney, fetal liver, thyroid, adrenal gland, stomach, small intestine, colon, muscle, heart, uterus, placenta, mammary gland, and testis cells. The described sequences were compiled from gene trapped cDNAs and clones isolated from a human testis cDNA library, and a human placenta cDNA (Edge Biosystems, Gaithersburg, MD). The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described genes, including the specifically described NHPs, and the NHP products; (b) nucleotides that encode one or more portions of the NHPs that correspond to functional domains, and the polypeptide products specified by such nucleotide sequences, including but not limited to the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHPs in which all or a part of at least one domain is deleted or altered, and the polypeptide products specified by such nucleotide sequences, including but not limited to soluble proteins and peptides in which all or a portion of the signal sequence is deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of an NHP, or one of its domains (e.g., a receptor binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or diagnostic derivatives of the described polynucleotides such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs comprising a sequence first disclosed in the Sequence Listing.

As discussed above, the present invention includes: (a) the human DNA sequences presented in the Sequence Listing (and vectors comprising the same) and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel F.M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at p. 2.10.3) and encodes a functionally equivalent gene product. Additionally contemplated are any nucleotide sequences that hybridize to the complement of the DNA sequence that encode and express an amino acid sequence presented in the Sequence Listing under moderately stringent conditions, e.g., washing in 0.2xSSC/0.1% SDS at 42°C (Ausubel et al., 1989, *supra*), yet still encode a functionally equivalent NHP product. Functional equivalents of a NHP include naturally occurring NHPs present in other species and mutant NHPs whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent No. 5,837,458). The invention also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding NHP ORFs, or their functional equivalents, encoded by polynucleotide sequences that are about 99, 95, 90, or about 85 percent similar to corresponding regions of SEQ ID NO:1 (as measured by BLAST sequence comparison analysis using, for example, the GCG sequence analysis package using standard default settings).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP gene nucleotide sequences. Such

hybridization conditions may be highly stringent or less highly stringent, as described above. In instances where the nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such molecules are generally about 16 to about 100 bases long, or about 20 to about 80, or about 34 to about 45 bases long, or any variation or combination of sizes represented therein that incorporate a contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc..

Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a micro array or high-throughput "chip" format). Additionally, a series of the described NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length may partially overlap each other and/or the NHP sequence may be represented using oligonucleotides that do not overlap. Accordingly, the described NHP polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 18, and preferably about 25, nucleotides in length that are each first disclosed in the described Sequence Listing. Such oligonucleotide sequences may begin at any nucleotide present within a sequence in the Sequence Listing and proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for

20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP gene antisense molecules, useful, for example, in NHP gene regulation (for and/or as antisense primers in amplification reactions of NHP gene nucleic acid sequences). With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further, such sequences may be used as part of ribozyme and/or triple helix sequences that are also useful for NHP gene regulation.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl)uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate,

a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier *et al.*, 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue *et al.*, 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue *et al.*, 1987, FEBS Lett. 215:327-330). Alternatively, double stranded RNA can be used to disrupt the expression and function of a targeted NHP.

Oligonucleotides of the invention can be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides can be synthesized by the method of Stein *et al.* (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin *et al.*, 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook *et al.*, 1989, Molecular Cloning, A Laboratory Manual (and periodic updates thereof), Cold Springs Harbor Press, N.Y.; and Ausubel *et al.*, 1989, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y.

Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to, nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), etc., that can be used in diagnostics and pharmacogenomics.

Further, a NHP gene homolog can be isolated from nucleic acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of mRNA prepared from, for example, human or non-human cell lines or tissue, such as prostate, rectum, colon, or adrenal gland, known or suspected to express an allele of a NHP gene.

The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the desired NHP gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via the screening of a genomic library.

PCR technology can also be used to isolate full length cDNA sequences. For example, RNA can be isolated, following standard

procedures, from an appropriate cellular or tissue source (i.e., one known, or suspected, to express a NHP gene, such as, for example, testis tissue). A reverse transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can be used, see e.g., Sambrook et al., 1989, *supra*.

A cDNA encoding a mutant NHP gene can be isolated, for example, by using PCR. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in an individual putatively carrying a mutant NHP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from an individual suspected of or known to carry a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, obesity, high blood pressure, etc.), or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express a mutant NHP allele. A

normal NHP gene, or any suitable fragment thereof, can then be labeled and used as a probe to identify the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP gene sequences can then be purified and subjected to sequence analysis according to methods well known to those skilled in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant NHP allele in an individual suspected of or known to carry such a mutant allele.

In this manner, gene products made by the putatively mutant tissue may be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.)

Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, AP-NHP or NHP-AP fusion proteins. In cases where a NHP mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to a NHP are likely to cross-react with a corresponding mutant NHP gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known in the art.

The invention also encompasses (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (*i.e.*, antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences (for example, baculo virus as described in U.S. Patent No. 5,869,336 herein incorporated by reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding

sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d) genetically engineered host cells that express an endogenous NHP gene under the control of an exogenously introduced regulatory element (*i.e.*, gene activation). As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the cytomegalovirus hCMV immediate early gene, regulatable, viral (particularly retroviral LTR promoters) the early or late promoters of SV40 adenovirus, the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast α -mating factors.

The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of the NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP gene (transcription factor inhibitors, antisense and ribozyme molecules, or gene or regulatory sequence replacement constructs), or promote the expression of a NHP (*e.g.*, expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, etc.).

The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs or inappropriately expressed NHPs for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for screening for drugs (or high throughput screening of combinatorial libraries)

effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of NHP in the body. The use of engineered host cells and/or animals may offer an advantage in that such systems allow not only for the
5 identification of compounds that bind to the endogenous receptor for an NHP, but can also identify compounds that trigger NHP-mediated signal transduction.

Finally, the NHP products can be used as therapeutics. For example, soluble derivatives such as NHP peptides/domains
10 corresponding the NHPs, NHP fusion protein products (especially NHP-Ig fusion proteins, *i.e.*, fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists (including compounds that modulate signal transduction which may act on
15 downstream targets in a NHP-mediated signal transduction pathway) can be used to directly treat diseases or disorders. For instance, the administration of an effective amount of soluble NHP, or a NHP-IgFc fusion protein or an anti-idiotypic antibody (or its Fab) that mimics the NHP could activate or effectively
20 antagonize the endogenous NHP receptor. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of a NHP, a NHP peptide, or a NHP fusion
25 protein to the body. Nucleotide constructs encoding functional NHPs, mutant NHPs, as well as antisense and ribozyme molecules can also be used in "gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological
30 disorders.

A knockout ES cell clone has been produced in a murine gene encoding an ortholog of the disclosed NHPs.

Various aspects of the invention are described in greater detail in the subsections below.

5.1 THE NHP SEQUENCES

5 The cDNA sequences (SEQ ID NOS: 1 and 3) and the
corresponding deduced amino acid sequence (SEQ ID NO: 2) of the
described NHPs are presented in the Sequence Listing. The NHP
genes were obtained from human testis and placenta cDNA libraries
using probes and/or primers generated from human gene trapped
10 sequence tags. Expression analysis has provided evidence that the
described NHPs can be expressed, for example, in human testis,
prostate, and gene trapped human cells. In addition to the genes
encoding trypsin inhibitors, the described NHPs share significant
similarity to a variety of cancer pathogenesis proteins, sperm
15 glycoproteins, and secretory proteins.

 The described open reading frames can also contain several
polymorphisms including an C to T transition corresponding to base
81 of SEQ ID NO:1, a G to C transversion corresponding to base 965
of SEQ ID NO:1 (changing a serine to a threonine), and a C to G
20 transversion corresponding to base 165 of the 5' UTR of SEQ ID
NO:3. SEQ ID NO:3 describes a full length ORF with flanking 5'
and 3' sequences.

5.2 NHPS AND NHP POLYPEPTIDES

25 NHPs, polypeptides, peptide fragments, mutated, truncated, or
deleted forms of the NHPs, and/or NHP fusion proteins can be
prepared for a variety of uses. These uses include but are not
limited to the generation of antibodies, as reagents in diagnostic
assays, the identification of other cellular gene products related
30 to a NHP, as reagents in assays for screening for compounds that
can be used as pharmaceutical reagents useful in the therapeutic
treatment of mental, biological, or medical disorders and disease.

002250" 08E'9960

The Sequence Listing discloses the amino acid sequences encoded by the described NHP genes. The NHPs have initiator methionines in DNA sequence contexts consistent with a translation initiation site, and further incorporate a hydrophobic leader sequence characteristic of secreted proteins.

The NHP amino acid sequences of the invention include the amino acid sequence presented in the Sequence Listing as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by the invention.

In fact, any NHP protein encoded by the NHP nucleotide sequences described above are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of the genetic code is well known, and, accordingly, each amino acid presented in the Sequence Listing, is generically representative of the well known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell *et al.* eds., Scientific American Books, New York, NY, herein incorporated by reference) are generically representative of all the various permutations and combinations of nucleic acid sequences that can encode such amino acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences as judged by any of a number of criteria, including, but not limited to, the ability to bind and cleave a substrate of a NHP, or the ability to effect an identical or complementary downstream signal transduction pathway, or a change in cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, etc.). Such functionally equivalent NHP

proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described above, but which result in a silent change, thus producing a functionally equivalent gene product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where, as in the present instance, the NHP peptide or polypeptide is thought to be a soluble or secreted molecule, the peptide or polypeptide can be recovered from the culture media. Such expression systems also encompass engineered host cells that express a NHP, or functional equivalent, *in situ*, i.e. anchored to the cell membrane. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well known to those skilled in the art. Alternatively, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of the NHP, but to assess biological activity, e.g., in drug screening assays.

The expression systems that may be used for purposes of the invention include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors

002260 08E 79660

containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*,
Pichia) transformed with recombinant yeast expression vectors
containing NHP nucleotide sequences; insect cell systems infected
with recombinant virus expression vectors (e.g., baculovirus)

5 containing NHP sequences; plant cell systems infected with
recombinant virus expression vectors (e.g., cauliflower mosaic
virus, CaMV; tobacco mosaic virus, TMV) or transformed with
recombinant plasmid expression vectors (e.g., Ti plasmid)
containing NHP nucleotide sequences; or mammalian cell systems
10 (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression
constructs containing promoters derived from the genome of
mammalian cells (e.g., metallothionein promoter) or from mammalian
viruses (e.g., the adenovirus late promoter; the vaccinia virus
7.5K promoter).

15 In bacterial systems, a number of expression vectors may be
advantageously selected depending upon the use intended for the
NHP product being expressed. For example, when a large quantity
of such a protein is to be produced for the generation of
pharmaceutical compositions of or containing NHP, or for raising
20 antibodies to a NHP, vectors that direct the expression of high
levels of fusion protein products that are readily purified may be
desirable. Such vectors include, but are not limited, to the *E.*
coli expression vector pUR278 (Ruther et al., 1983, EMBO J.
2:1791), in which a NHP coding sequence may be ligated
25 individually into the vector in frame with the *lacZ* coding region
so that a fusion protein is produced; pIN vectors (Inouye &
Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke &
Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX
vectors may also be used to express foreign polypeptides as fusion
30 proteins with glutathione S-transferase (GST). In general, such
fusion proteins are soluble and can easily be purified from lysed
cells by adsorption to glutathione-agarose beads followed by

elution in the presence of free glutathione. The PGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

5 In an insect system, *Autographa californica* nuclear polyhydrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. A NHP gene coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed
10 under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of NHP gene coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These
15 recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed (e.g., see Smith et al., 1983, J. Virol. 46: 584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as
20 an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-
25 essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., See Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific
30 initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression

002260" 08E/9960

vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (See Bittner et al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the NHP sequences described above can be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with

002250" 08E'9950

DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be
5 allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into
10 cell lines. This method may advantageously be used to engineer cell lines which express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

A number of selection systems may be used, including but not
15 limited to the herpes simplex virus thymidine kinase (Wigler, et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy, et al., 1980, Cell 22:817) genes
20 can be employed in tk⁻, hgp⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler, et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare, et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt,
25 which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, et al., 1981, J. Mol. Biol. 150:1); and hyg^r, which confers resistance to hygromycin (Santerre, et al., 1984, Gene 30:147).

30 Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described by Janknecht et al.

allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht, et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺-nitriloacetic acid-agarose columns and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

5.3 ANTIBODIES TO NHP PRODUCTS

Antibodies that specifically recognize one or more epitopes of a NHP, or epitopes of conserved variants of a NHP, or peptide fragments of a NHP are also encompassed by the invention. Such antibodies include but are not limited to polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

The antibodies of the invention may be used, for example, in the detection of NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes, as described, below, in Section 5.5, for the evaluation of the effect of test compounds on expression and/or activity of a NHP gene product. Additionally, such antibodies can be used in conjunction gene therapy to, for example, evaluate the normal and/or engineered NHP-expressing cells prior to their introduction into the patient. Such antibodies may additionally be used as a method for the inhibition

of abnormal NHP activity. Thus, such antibodies may, therefore, be utilized as part of treatment methods.

For the production of antibodies, various host animals may be immunized by injection with the NHP, an NHP peptide (e.g., one corresponding to a functional domain of an NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of the NHP or mutated variant of the NHP. Such host animals may include but are not limited to pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid, ovalbumin, cholera toxoid or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, can be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Patent No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal

Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be
5 cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, Proc. Natl.

10 Acad. Sci., 81:6851-6855; Neuberger *et al.*, 1984, Nature, 312:604-608; Takeda *et al.*, 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity
15 can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region. Such technologies are described in U.S. Patents Nos. 6,075,181
20 and 5,877,397 and their respective disclosures which are herein incorporated by reference in their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent 4,946,778; Bird, 1988, Science 242:423-426; Huston *et al.*, 1988, Proc. Natl.

25 Acad. Sci. USA 85:5879-5883; and Ward *et al.*, 1989, Nature 334:544-546) can be adapted to produce single chain antibodies against NHP gene products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

30 Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the

002260" DBE 29950

Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of

5 monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" a given NHP, using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, 1993, FASEB J 7(5):437-444; and Nissinoff, 1991, 10 J. Immunol. 147(8):2429-2438). For example antibodies which bind to a NHP domain and competitively inhibit the binding of NHP to its cognate receptor can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a receptor. Such anti-idiotypic antibodies or fragments of such 15 anti-idiotypes can be used in therapeutic regimens involving a NHP signaling pathway.

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and 20 functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the 25 scope of the appended claims.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising at least 24 contiguous bases of nucleotide sequence first disclosed
5 in the NHP gene described in SEQ ID NO: 1.

2. An isolated nucleic acid molecule comprising a nucleotide sequence that:

- 10 (a) encodes the amino acid sequence shown in SEQ ID NO: 2; and
- (b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO: 1 or the complement thereof.

15 3. An expression vector comprising a polynucleotide sequence encoding the amino acid sequence shown in SEQ ID NO:2.

[illegible]

25

PATENT APPLICATION

DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

ATTORNEY DOCKET NO. LEX-0042-USA

As a below named inventor, I hereby declare that:

My residence/post office address and citizenship are as stated below next to my name;

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Novel Human Protease Inhibitor-Like Proteins and Polynucleotides Encoding the Same

the specification of which is attached hereto unless the following box is checked:

☐ was filed on _____ as US Application Serial No. or PCT International Application

Number _____ and was amended on _____ (if applicable).

I hereby state that I have reviewed and understood the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above. I acknowledge the duty to disclose all information which is material to patentability as defined in 37 CFR 1.56.

Foreign Application(s) and/or Claim of Foreign Priority

I hereby claim foreign priority benefits under Title 35, United States Code Section 119 of any foreign application(s) for patent or inventor(s) certificate listed below and have also identified below any foreign application for patent or inventor(s) certificate having a filing date before that of the application on which priority is claimed:

COUNTRY	APPLICATION NUMBER	DATE FILED	PRIORITY CLAIMED UNDER 35 U.S.C. 119
			YES: _____ NO: _____
			YES: _____ NO: _____

Provisional Application

I hereby claim the benefit under Title 35, United States Code Section 119(e) of any United States provisional application(s) listed below:

APPLICATION SERIAL NUMBER	FILING DATE
60/156,101	09/24/1999

U.S. Priority Claim

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NUMBER	FILING DATE	STATUS(patented/pending/abandoned)

POWER OF ATTORNEY:

As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) listed below to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Lance K. Ishimoto, Reg. No. 41866

Send Correspondence to:

Lance K. Ishimoto
Lexicon Genetics Incorporated
4000 Research Forest Drive
The Woodlands, TX 77381

Direct Telephone Calls To:

Lance K. Ishimoto
(281) 362-6554

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION (continued)

ATTORNEY DOCKET NO. LEX-0042-USA

Full Name of Inventor: Gregory Donoho

Citizenship: USA

Residence: 41 Fallenstone Dr., The Woodlands, TX 77381

Post Office Address: Same

Inventor's Signature

Date

Full Name of Inventor: C. Alexander Turner, Jr.

Citizenship: USA

Residence: 67 Winter Wheat Place, The Woodlands, TX 77381

Post Office Address: Same

Inventor's Signature

Date

Full Name of Inventor: Frank Wattler

Citizenship: Germany

Residence: Bennostrasse 11a, Stockdorf, Germany 82131

Post Office Address: Same

Inventor's Signature

Date

Full Name of Inventor: Michael C. Nehls

Citizenship: Germany

Residence: Paul-Keller-Strasse 6, Stockdorf, Germany 82131

Post Office Address: Same

Inventor's Signature

Date

Full Name of Inventor: Glenn Friedrich

Citizenship: Canada

Residence: c/o Breland & Breland, Houston, TX 77004

Post Office Address: Same

Inventor's Signature

Date

Full Name of Inventor: Brian Zambrowicz

Citizenship: USA

Residence: 18 Firethorne Place, The Woodlands, TX 77382

Post Office Address: Same

Inventor's Signature

Date

Full Name of Inventor: Arthur T. Sands

Citizenship: USA

Residence: 163 Bristol Bend Circle, The Woodlands, TX 77382

Post Office Address: Same

Inventor's Signature _____

Date _____

002260 0324960

SEQUENCE LISTING

<110> Donoho, Gregory
Turner, C. Alexander Jr.
Wattler, Frank
Nehls, Michael
Friedrich, Glenn
Zambrowicz, Brian
Sands, Arthur T.

<120> Novel Human Protease Inhibitor-Like
Proteins and Polynucleotides Encoding the Same

<130> LEX-0042-USA

<150> US 60/156,101

<151> 1999-09-24

<160> 3

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1491

<212> DNA

<213> homo sapiens

<400> 1
atgagctgcg tcctggggtgg tgtcatcccc ttggggctgc tgttcctggt ctgcgggatcc 60
caaggctacc tcctgcccac cgctactctc ttagaggagc tgctcagcaa ataccagcac 120
aacgagtctc actcccggtt ccgcagagcc atccccaggg aggacaagga ggagatcctc 180
atgctgcaca acaagcttcg gggccaggtg cagcctcagg cctccaacat ggagtacatg 240
acctgggatg acgaactgga gaagtctgct gcagcgtggg ccagtcagtg catctgggag 300
cacgggcccc ccagtctgct ggtgtccatc gggcagaacc tgggcgctca ctggggcagg 360
tategtctc cggggttcca tgtgcagtc tggtatgac aggtgaagga ctacacctac 420
ccctaccgca gcgagtgcaa cccctggtgt ccagagaggt gctcggggcc tatgtgcacg 480
cactacacac agatagtttg ggccaccacc aacaagatcg gttgtgctgt gaacacctgc 540
cggaagatga ctgtctgggg agaagtttgg gagaacgcgg tctactttgt ctgcaattat 600
tctccaaagg ggaactggat tggagaagcc ccctacaaga atggccggcc ctgctctgag 660
tgccccacca gctatggagg cagctgcagg aacaacttgt gttaccgaga agaaacctac 720
actccaaaac ctgaaacgga cgagatgaat gaggtggaaa cggtcccat tcctgaagaa 780
aaccatgttt ggctccaacc gaggtgatg agaccacca agcccaagaa aacctctgcg 840
gtcaactaca tgacccaagt cgctcagatgt gacaccaaga tgaaggacag gtgcaaaggg 900
tccacgtgta acaggtacca gtgccagca ggctgcctga accacaaggc gaagatcttt 960
ggaagtctgt tctatgaaag ctctgttagc atatgccgag ccgccatcca ctacgggatc 1020
ctggatgaca agggaggcct ggtggatata accaggaacg ggaaggctcc cttcttcgtg 1080
aagtctgaga gacacggcgt gcagtcctc agcaaataca aaccttcag ctcatctatg 1140
gtgtcaaaaag tgaaagtgca ggatttggac tgctacacga ccgttgctca gctgtgccccg 1200
tttgaaaagc cagcaactca ctgcccaga atccattgtc cggcacactg caaagacgaa 1260
ccttcctact gggctccggt gtttggaaac aacatctatg cagatacctc aagcatctgc 1320
aagacagctg tgcacgcggg agtcatcagc aacgagagtg ggggtgacgt ggacgtgatg 1380
cccgtggata aaaagaagac ctacgtgggc tcgctcagga atggagttca gtctgaaagc 1440
ctggggactc ctcgggatgg aaaggccttc cggatctttg ctgtcaggca g 1491

<210> 2

<211> 497

<212> PRT

<213> homo sapiens

<400> 2

Met Ser Cys Val Leu Gly Gly Val Ile Pro Leu Gly Leu Leu Phe Leu

002260" 08E/9960

1	5	10	15
Val Cys Gly Ser Gln Gly Tyr Leu Leu Pro Asn Val Thr Leu Leu Glu			
20	25	30	
Glu Leu Leu Ser Lys Tyr Gln His Asn Glu Ser His Ser Arg Val Arg			
35	40	45	
Arg Ala Ile Pro Arg Glu Asp Lys Glu Glu Ile Leu Met Leu His Asn			
50	55	60	
Lys Leu Arg Gly Gln Val Gln Pro Gln Ala Ser Asn Met Glu Tyr Met			
65	70	75	80
Thr Trp Asp Asp Glu Leu Glu Lys Ser Ala Ala Ala Trp Ala Ser Gln			
85	90	95	
Cys Ile Trp Glu His Gly Pro Thr Ser Leu Leu Val Ser Ile Gly Gln			
100	105	110	
Asn Leu Gly Ala His Trp Gly Arg Tyr Arg Ser Pro Gly Phe His Val			
115	120	125	
Gln Ser Trp Tyr Asp Glu Val Lys Asp Tyr Thr Tyr Pro Tyr Pro Ser			
130	135	140	
Glu Cys Asn Pro Trp Cys Pro Glu Arg Cys Ser Gly Pro Met Cys Thr			
145	150	155	160
His Tyr Thr Gln Ile Val Trp Ala Thr Thr Asn Lys Ile Gly Cys Ala			
165	170	175	
Val Asn Thr Cys Arg Lys Met Thr Val Trp Gly Glu Val Trp Glu Asn			
180	185	190	
Ala Val Tyr Phe Val Cys Asn Tyr Ser Pro Lys Gly Asn Trp Ile Gly			
195	200	205	
Glu Ala Pro Tyr Lys Asn Gly Arg Pro Cys Ser Glu Cys Pro Pro Ser			
210	215	220	
Tyr Gly Gly Ser Cys Arg Asn Asn Leu Cys Tyr Arg Glu Glu Thr Tyr			
225	230	235	240
Thr Pro Lys Pro Glu Thr Asp Glu Met Asn Glu Val Glu Thr Ala Pro			
245	250	255	
Ile Pro Glu Glu Asn His Val Trp Leu Gln Pro Arg Val Met Arg Pro			
260	265	270	
Thr Lys Pro Lys Lys Thr Ser Ala Val Asn Tyr Met Thr Gln Val Val			
275	280	285	
Arg Cys Asp Thr Lys Met Lys Asp Arg Cys Lys Gly Ser Thr Cys Asn			
290	295	300	
Arg Tyr Gln Cys Pro Ala Gly Cys Leu Asn His Lys Ala Lys Ile Phe			
305	310	315	320
Gly Ser Leu Phe Tyr Glu Ser Ser Ser Ser Ile Cys Arg Ala Ala Ile			
325	330	335	
His Tyr Gly Ile Leu Asp Asp Lys Gly Leu Val Asp Ile Thr Arg			
340	345	350	
Asn Gly Lys Val Pro Phe Phe Val Lys Ser Glu Arg His Gly Val Gln			
355	360	365	
Ser Leu Ser Lys Tyr Lys Pro Ser Ser Ser Phe Met Val Ser Lys Val			
370	375	380	
Lys Val Gln Asp Leu Asp Cys Tyr Thr Thr Val Ala Gln Leu Cys Pro			
385	390	395	400
Phe Glu Lys Pro Ala Thr His Cys Pro Arg Ile His Cys Pro Ala His			
405	410	415	
Cys Lys Asp Glu Pro Ser Tyr Trp Ala Pro Val Phe Gly Thr Asn Ile			
420	425	430	
Tyr Ala Asp Thr Ser Ser Ile Cys Lys Thr Ala Val His Ala Gly Val			
435	440	445	
Ile Ser Asn Glu Ser Gly Gly Asp Val Asp Val Met Pro Val Asp Lys			
450	455	460	
Lys Lys Thr Tyr Val Gly Ser Leu Arg Asn Gly Val Gln Ser Glu Ser			
465	470	475	480
Leu Gly Thr Pro Arg Asp Gly Lys Ala Phe Arg Ile Phe Ala Val Arg			
485	490	495	
Gln			

<210> 3
 <211> 2272
 <212> DNA
 <213> homo sapiens

<400> 3
 cccagggcgt ctccggctgc tcccattgag ctgtctgctc gctgtgcccg ctgtgcctgc 60
 tgtgcccgcg ctgtgcgcgc tgctaccgcg tctgtctggac gcgggagacg ccagcgagct 120
 ggtgattgga gccctgcgga gagctcaagc gccagctct gcccaggag cccaggtgc 180
 cccgtgagtc ccatagttgc tgcaggagtg gagccatgag ctgctcctg ggtggtgtca 240
 tccccttggg gctgctgttc ctgggtctgc gatcccaagg ctacctctg cccaacgtca 300
 ctctcttaga ggagctgctc agcaaatacc agcacaacga gtctcactcc cgggtccgca 360
 gagccatccc cagggaggac aaggaggaga tcctcatgct gcacaacaag ctccggggcc 420
 aggtgcagcc tcaggcctcc aacatggagt acatgacctg ggatgacgaa ctggagaagt 480
 ctgctgcagc gtgggccagt cagtgcattc gggagcacgg gccaccagt ctgctggtgt 540
 ccatcgggca gaacctgggc gctcactggg gcaggtatcg ctctccgggg ttccatgtgc 600
 agtcctggta tgacgaggtg aaggactaca cctaccctta cccgagcagag tgcaaccctt 660
 ggtgtccaga gaggtgctcg gggcctatgt gcacgcacta cacacagata gtttgggcca 720
 ccaccaacaa gatcggttgt gctgtgaaca cctgccggaa gatgactgtc tggggagaag 780
 tttgggagaa cgcggtctac tttgtctgca attattctcc aaaggggaac tggattggag 840
 aagcccccta caagaatggc cggccctgct ctgagtgcgc acccagctat ggaggcagct 900
 gcaggaacaa cttgtgttac cgagaagaaa cctacactcc aaaacctgaa acggacgaga 960
 tgaatgaggt ggaaacggct cccattcctg aagaaaacca tgtttggctc caaccgaggg 1020
 tgatgagacc caccaagccc aagaaaacct ctgcggtcaa ctacatgacc caagtcgtca 1080
 gatgtgacac caagatgaag gacaggtgca aagggtccac gtgtaacagg taccagtgc 1140
 cagcaggctg cctgaaccac aaggcgaaga tctttggaag tctgttctat gaaagctcgt 1200
 ctagcatatg ccgcgccgcc atccactacg ggatcctgga tgacaaggga ggcctggtgg 1260
 atatcaccag gaacgggaag gtccccttct tcgtgaagtc tgagagacac ggcgtgcagt 1320
 ccctcagcaa atacaaacct tccagctcat tcatggtgtc aaaagtgaag gtgcaggatt 1380
 tggactgcta cagcaccgtt gctcagctgt gccggttga aaagccagca actcactgcc 1440
 caagaatcca ttgtccggca cactgcaaag acgaaccttc ctactgggct ccggtgtttg 1500
 gaaccaacat ctatgcagat acctcaagca tctgcaagac agctgtgcac gcgggagtca 1560
 tcagcaacga gagtgggggt gacgtggacg tgatgccgt ggataaaaag aagacctacg 1620
 tgggctcgct caggaatgga gttagctctg aaagcctggg gactcctcgg gatggaaagg 1680
 ccttcgggat ctttgcctgc aggcagtga tttccagcac caggggagaa ggggcgtctt 1740
 caggaggggt tcgggggttt gcttttattt ttattttgtc attgcgggggt atatggagag 1800
 tcaggaaact tcctttgact gatgttcagt gtccatcact ttgtggcctg tgggtgaggt 1860
 gacatctcat cccctcactg aagcaacagc atcccaaggt gctcagccgg actccctggt 1920
 gcctgatcct gctggggcct gggggtctcc atctggacgt cctctctcct ttagagatct 1980
 gagctgtctc ttaaagggga cagttgccca aaatgttctc tgctatgtgt tcttctgttg 2040
 gtggaggaag ttgatttcaa cccccctgcc aaaagaacaa accatttgaa gctcacaatt 2100
 gtgaagcatt cacggcgtcg gaagaggcct tttgagcaag cgccaatgag ttccaggaat 2160
 gaagtagaag gtatgtattt aaaaataaaa aacacagtcc gtccctacca atagaggaaa 2220
 atgggttttaa tgtttgctgg tcagacagac aaatgggcta gagtaagaag gc 2272